

What Is Abortion Pill And How To Take It?

Early Pregnancy Termination With Medical Abortion Method


<https://abortionpillrx247.com/>

SAM ADAM Dallas, Texas, USA

ABORT YOUR PREGNANCY SAFELY WITH MTP KIT



COMPOSITION OF ABORTION PILL



Order Abortion Pill online which holds 5 pills combo pack.

1 Pills of Mifepristone 200 mg

4 tablets of Misoprostol 200 mcg (each pill holds 200 mcg dosages power)

ABORTION PILL DOSAGE FORM

Mifepristone Pill must be taken orally with a glass of water after intake of food.

Misoprostol tablets can be taken buccally or vaginally as per your choice.

PURPOSE TO USE

[Buy Abortion Pills online](#) which is used for the termination of pregnancy at home with the self-induced abortion method.

WHEN TO USE?


From starting days of gestation to 9 weeks, i.e. still 63 days of gestation.

THE AVAILABLE BRAND OF ABORTION PILL

MTP Kit, Mifegest Kit, and Mifeprex Kit

WORKING OF ABORTION PILL

Purchase Abortion Pill online which contains Mifepristone and Misoprostol pill combo, and each pill works differently to terminate pregnancy.



MIFEPRISTONE

Order Mifepristone online which to prevent fetal particle growth. Also helps to get out of the pregnancy tissue from the uterus.

MISOPROSTOL

Buy Misoprostol online which helps to expand the cervix and tighten the uterus. So, it can help rinse out the parts of the pregnancy from the uterus.

HOW TO TAKE THE ABORTION PILL?



MIFEPRISTONE

Buy Mifepristone online which is called an antiprogestin pill. This is the primary taken medicine, which must have to take 1st when you decided on pregnancy termination. This must be taken orally with a glass of water after having a meal.

MISOPROSTOL PROSTAGLANDIN PILL

Order Misoprostol online which is called a prostaglandin pill. This is secondary medicine, which must have taken after a 24-hour intake of Mifepristone 200mg. You must have to take 2-2 dosages of Misoprostol Abortion Pill at a time with 6/12/24 hour gaps.

There are two ways in which can be taken this pill, i.e. buccally or vaginally




BUCCALLY


Put 2-2 pills at a time in the cheek. Don't crush or chew it. Keep at least 30 minutes of this medicine in the cheek to get dissolved in the mouth properly. After completing 30 minutes if this pregnancy termination pill remains in then swallow it with a glass of water.

VAGINALLY

Put all 4 tablets at a time vaginally in a sequence.



WHAT ARE THE SIDE EFFECTS?



There is no major side effect you experience after inducing it. There is some minor side effect which also gets disappeared within 2 weeks of completing the pregnancy termination process.

Nausea, Vomiting, Diarrhea, Dizziness, Fever, Stomach cramp, Uterine contractions, Menorrhagia (heavy menstrual bleeding)